

DEFOAMER

LIQUID DEFOAM CONCENTRATE

**USE WHEN
FOAM IS A
PROBLEM**

**Whenever and wherever foam is
a problem:**

**Carpet shampoo pickup
Sewage systems
Decorative water fountains
Whirlpools**

**No solvents, caustics,
acids or fumes**

**QUICKLY
PENETRATES
AND BREAKS
DOWN BUBBLES**

**Ready to use product.
Helps prevent foam from
clogging vacuums.
Keeps fountains from
becoming a nuisance.**

STOP FOAM

PCC PRODUCTS

6400 Herman Avenue • Cleveland, Ohio 44102

216-281-1155 • 800-233-5078

FAX 216-281-1186

www.prod-chem.com

DESCRIPTION

This concentrated product may be used anywhere that suds cause a problem. A defoamer silicone oil properly emulsified in water to allow the silicone oil to float-out on the surface of the foaming liquid and cancel the foam quickly.

APPLICATION

For use in the pickup tank of steam or hot water extractors.
For use in sewage treatment plants, decorative water fountains and whirlpools.

DIRECTIONS FOR USE

Prior to using wet recovery vacuums:

1. With unit turned on, suck up a small amount of DEFOAMER to coat hose lining. Then,
2. Pour 2 ounces DEFOAMER into recovery tank.

Defoamer may be used full strength anywhere foam is a problem in such places as water treatment applications, sewers, and drains.

SPECIFICATIONS

Packaged: 5- gallon pail ; 4- 1 gallon case
Appearance: Milky liquid
Fragrance: Characteristic
pH: 7.50
Dilution: Use as necessary until foam subsides
Flash point: None. Non-flammable
Solubility in water: Complete
Authorization: USDA authorized for use in federally inspected meat and poultry plants

PRECAUTIONS

Keep out of reach of children. Possible eye or skin irritant. Do not get in eyes, on skin, or on clothing. Rinse empty container with water and discard. This product is intended for industrial and commercial use and storage by trained service persons only. Manufacturer will not be responsible for any injury, loss, or damage, if product is used in any manner not in compliance with label directions or if precautions are not observed.

FIRST AID

EYES: Flush with large amounts of water for at least 15 minutes; contact physician. IF SWALLOWED: Induce vomiting. Drink large quantities of water. Consult physician. EXTERNAL: Flush with large amounts of water.